

County Environmental Health Profiles Sedgwick County, Kansas

Exposure: Outdoor Air

Air pollution means that there are substances in the air that should not be there, or should be there in smaller amounts. Two important pollutants to consider for the health of a county are fine particulate matter and ozone. Particulate matter describes particles that settle in the lungs after being inhaled. Particulate matter 2.5 (PM_{2.5}) refers to the size of the particles. Ozone is created as a result of emissions from vehicles and industrial facilities interacting with sunlight. Both particulate matter and ozone can trigger health problems, especially within children, the elderly, and individuals with lung disease such as asthma. The United States Environmental Protection Agency sets National Ambient Air Quality Standards (NAAQS) for pollutants considered harmful to public health and the environment.





Exposure: Indoor Air: Radon

Radon is a cancer-causing radioactive gas. It is tasteless, odorless, colorless, and produced naturally from the decay of uranium found in nearly all soils. Radon gas moves from the ground under and around homes through cracks and other holes in the foundation. The United States Environmental Protection Agency recommends corrective action be taken if the radon level in a home is 4 picocuries per liter (pCi/L) or higher. The only way to know what the radon level is in your home is to test.

The table below shows the average annual radon level in Sedgwick County homes in 2010 to be **3.6 pCi/L**, which is below the state average of **4.8 pCi/L**. The annual percent of radon measurements at or above 4 pCi/L for 2010 was also less than the state average. Sedgwick County's percent at or above the suggested action level was **30.9%** and the Kansas average was **41.0%**.

) _ u! _ d		
Period	Sedgwick County	Kansas
2010	3.6	4.8
	C.1.C.U.	

Annual maximum radon level reported (pCi/L).	2010	92.8	260
Annual total number of radon measurements.	2010	3,771	50,182
Annual total number of radon measurements at or above 4 pCi/L.	2010	1,167	20,592
Annual percent of radon measurements at or above 4 pCi/L.	2010	30.9	41.0
Source: Kansas Radon Program			

Exposure: Age of Housing

Lead is found throughout our environment. It is a naturally occurring bluish-gray metal found in small amounts in the Earth's crust. A good amount of lead in our environment comes from human activities including burning fossil fuels, mining, and manufacturing. In the United States, the most common source of exposure for lead-poisoned children is lead-based paint. Homes built before 1980 represent a potential source for lead exposure because they likely contain lead-based paint. The vast majority of homes built before 1950 contain substantial amounts of lead-based paint. The health effects of lead exposure include intellectual and behavioral deficit in children.

In 2010, **18.8%** of homes in Sedgwick County were built before 1950, compared to **25.1%** in Kansas. In 2010, the percent of homes built between 1950 and 1979 in Sedgwick County was **43.0%**, which was higher than the **40.0%** statewide.

Exposure: Age of Housing			
Indicator	Period	Sedgwick County	Kansas
Number of homes built before 1950	2010	39,316	307,414
Number of homes built between 1950 and 1979	2010	89,706	489,045
Percent of homes built before 1950	2010	18.8	25.1
Percent of homes built between 1950 and 1979	2010	43.0	40.0
Source: U.S. Census Bureau			

Health Outcome: Childhood Blood Lead Poisoning

Lead poisoning is a preventable public health problem, especially in children. The United States Centers for Disease Control and Prevention now recommends that case management occur for any child with a blood lead level of 5 or more micrograms per deciliter (5 ug/dL). However, several studies have shown that there is no safe level for blood lead poisoning. Therefore, it is recommended that all children be tested for blood lead by the age of 3. Of all the children born in 2010 in Sedgwick County, **10.1%** were tested before age 3, compared to the **16.9%** of all children born in Kansas in the same year.

Health Outcome: Childhood Blood Lead			
Indicator	Period	Sedgwick County	Kansas
Percent of children born in the same year and tested before age 3	2010 cohort	10.1	16.9
Percent of children tested during the period with confirmed blood lead levels of 5 µg/dL or greater	2012	1.5	2.1
Source: Kansas Environmental Public Health Tracking Program, Kansas Department of Health and Environment			

Health Outcome: Carbon Monoxide Poisoning

Carbon monoxide (CO) is an odorless, colorless, tasteless gas which can cause sudden illness and even death. Carbon monoxide is produced when a fossil fuel is not burned completely. Carbon monoxide levels in the home are created by a number of different sources, including kerosene and gas space heaters, furnaces, wood stoves, generators, and other gasoline-powered equipment. Tobacco smoke also contributes to indoor CO levels.

Due to the small number of hospitalizations from carbon monoxide poisoning, the data for Sedgwick County cannot be displayed.

Health Outcome: Carbon Monoxide Hospitalization			
Indicator	Period	Sedgwick County	Kansas

Age-adjusted rate of hospitalization for CO poisoning per 100,000 population (all causes included)	2012	N/A	0.55
Number of hospitalizations for CO poisoning	2008-2012	Suppressed*	122

Source: Kansas hospital discharge data. Counts for the year are based on date of admission.

Health Outcome: Asthma

Asthma is a chronic condition that affects the airways that carry air in and out of a person's lungs. If a person has asthma, the inside of these airways is irritated and swollen. Some people with asthma experience asthma attacks. In that case, symptoms only appear when they are exposed to something that irritates their breathing. Others have a kind of asthma that makes breathing difficult all of the time.

Sedgwick County had an age-adjusted rate of **9.9** hospitalizations for asthma per 10,000 population for the year 2012. This rate is above the state average of **8.5** hospitalizations per 10,000 population.

Health Outcome: Asthma			
Indicator	Period	Sedgwick County	Kansas
Age-adjusted rate of hospitalization for asthma per 10,000 population. [‡]	2012	9.9	8.5
Percent of adults ever diagnosed with asthma.§	2011	N/A	12.9
Percent of adults with current asthma. §§	2013	8.8	8.9
Percent of adults with active asthma with an attack in the past year. §	2011	N/A	54.1
Percent of children ever diagnosed with asthma. §	2012	N/A	13.9

[‡] Kansas hospital discharge data. Note: Counts for the year are based on date of admission.

^{*} The data is suppressed to protect the privacy of the persons affected

N/A= Statistic is "Not Available" due to very small number of events

[§] Kansas Behavioral Risk Factor Surveillance Survey.

 $^{^{\}S\S}$ Kansas Behavioral Risk Factor Surveillance Survey (local Data). N/A= Statistic is "Not Available" due to very small number of events

Health Outcome: Reproductive and Birth Outcomes

Reproduction is complex, and many factors affect a mother's ability to conceive, carry a baby to term, and deliver a baby without complications. Environmental factors such as air pollution and exposure to contaminated drinking water can increase the likelihood of low birth weight and preterm births.

The Total Fertility Rate is a measure that controls for the fact that a woman's fertility (her ability to bear children) varies over time and space. It is calculated for a cohort of 1,000 women. It is a more precise measure than the general fertility rate and allows comparison between regions and age groups.

During the 2009 to 2013 timeframe, Sedgwick County had 7.7 infant deaths per 1,000 live births. This rate was higher than the statewide rate during the same time period. From 2011-2013, Sedgwick County was above the state average for the percent of low birthweight for live, term, singleton births with 8.1% versus the state average of 7.1%. In 2010 the total fertility rate for Sedgwick County was **2,325.9** compared to the **2,148.8** Kansas rate.

Population: Reproductive and Bir Indicator	Period	Sedgwick County	Kansas
Infant mortality rate per 1,000 live births	2009-2013	7.7	6.4
Male to Female sex ratio§	2010	1.032	1.031
Total fertility rate per 1,000 women of reproductive age§	2010	2,325.9	2,148.8
Percent of low birthweight (less than 2500 grams) live term singleton births [‡]	2011-2013	8.1	7.1
[‡] Kansas Department of Health and Environment			
§ National Center for Health Statistics (NCHS). Term si	ngleton only		

Health Outcome: Birth Defects

A birth defect is a problem that happens while a baby is developing. Most birth defects happen during the first three months of pregnancy. Major congenital birth defects are conditions present at birth

causing structural changes in one or more parts of the body. A number of environmental exposures, including exposures to certain chemicals, can cause the development of birth defects.

From 2008-2012 Sedgwick County had a higher Prevalence of trisomy 21 per 10,000 live births to mothers 35 years old or older at delivery compared to the state. Kansas has a rate of **35.8** and Sedgwick County has a rage of **42.8**. All other data shows rates or below the state averages provided.

Health Effects: Birth Defects			
Indicator	Period	Sedgwick County	Kansas
Prevalence of anencephaly per 10,000 live births [‡]	2008-2012	Suppressed*	1.7
Prevalence of cleft lip with or without cleft palate per 10,000 live births [‡]	2008-2012	6.8	7.6
Prevalence of cleft palate without cleft lip per 10,000 live births [‡]	2008-2012	2.8	3.6
Prevalence of gastroschisis per 10,000 live births [‡]	2008-2012	4.2	4.2
Prevalence of hypospadias per 10,000 live births [‡]	2008-2012	6.5	10.8
Prevalence of trisomy 21 per 10,000 live births to mothers less than 35 years old at delivery [‡]	2008-2012	3.2	4.6
Prevalence of trisomy 21 per 10,000 live births to mothers 35 years old or older at delivery [‡]	2008-2012	42.8	35.8
* Kansas Department of Health and Environment * The data is suppressed to protect the privacy of the	persons affected		

Health Outcome: Cancer

Cancer is a diverse group of diseases characterized by the uncontrolled growth and spread of abnormal cells. Cancer is believed to be caused by both external and internal risk factors.

During the 2007 to 2011 timeframe, the age-adjusted incidence of thyroid cancer per 100,000 population for Sedgwick County was **18.7**, which was above the state average of **15.6**. Age-adjusted

incidence of liver and biliary tree cancer per 100,000 population for Sedgwick County was **8.5** which was above the state average of **7.3**. Age-adjusted incidence of lung and bronchus cancer per 100,000 population and Age-adjusted incidence of kidney and renal pelvis cancer per 100,000 population were slightly above the state rate. All other cancers listed were at or below the state average.

Health Effects: Selected Cancer Sit Indicator	Period	Sedgwick County	Kansas
Age-adjusted incidence of female breast cancer per 100,000 female population [‡]	2007-2011	117.4	122.1
Age-adjusted incidence of brain cancer per 100,000 population [‡]	2007-2011	6.0	6.2
Age-adjusted incidence of lung and bronchus cancer per 100,000 population [‡]	2007-2011	62.2	62.0
Age-adjusted incidence of thyroid cancer per 100,000 population [‡]	2007-2011	18.7	15.6
Age-adjusted incidence of kidney and renal pelvis cancer per 100,000 population [‡]	2007-2011	16.9	16.6
Age-adjusted incidence of bladder cancer (in situ included) per 100,000 population [‡]	2007-2011	19.7	22.0
Age-adjusted incidence of leukemia per 100,000 population [‡]	2007-2011	12.9	13.8
Age-adjusted incidence of liver and biliary tree cancer per 100,000 population [‡]	2007-2011	8.5	7.3
Age-adjusted incidence of melanoma of the skin per 100,000 population [‡]	2007-2011	21.9	23.0
Age-adjusted incidence of pancreatic cancer per 100,000 population [‡]	2007-2011	9.8	10.2
[‡] Kansas Department of Health and Environment [£] Unstable rate (case count less than 20) N/A= "Not Available" due to very small number of eve	ents		

Health Outcome: Acute Myocardial Infarction

Acute myocardial infarction, or heart attacks, occurs when the vessels supplying blood to the heart become blocked. The resulting blood and oxygen shortage causes damage and potential death of heart tissue. Several factors may increase the risks of heart attacks. These factors include health, lifestyle, and environmental factors. For example, air pollution has been linked to decreases in lung function and heart attacks. In addition, research has shown that high levels of air pollution may affect people with pre-existing conditions like asthma and heart disease.

In 2012, Sedgwick County had a rate of **32.7** hospitalizations per 10,000 population, which was less than the **26.3** hospitalizations per 10,000 population rate statewide.

Health Effects: Acute Myocardial Infarction (Heart Attack)			
Indicator	Period	Sedgwick County	Kansas
Age-adjusted rate of hospitalization for heart attack per 10,000 population among persons 35 years old and over. ‡	2012	32.7	26.3
[‡] Kansas hospital discharge data. Note: Counts for the year are based on date of admission.			

Population Characteristics

Population data, both the number of people and their location, are important for determining the impact of a disease on a community and society at large, and the trends and patterns of disease over space and time. Characteristics of the populations (sex, age, race, and ethnicity) and their socioeconomic measures (poverty) have been associated with disease incidence and prevalence.

In 2012, the median household income for Sedgwick County was \$47,649 which is \$2,491 less than the state median household income. During the same time period, a higher percentage of people living in Sedgwick County lived in poverty than the state average.

Population: Demographic & Socioeconomic Characteristics				
Indicator	Period	Sedgwick County	Kansas	
Estimated total population	2013	505,415	2,893,957	

2012	47,649	50,140
2000	85.1	86.0
2012	16.4	14.0
	2000	2000 85.1